

Bicherón 50NF

Super Concentrated Insecticide

Cypermethrin 20%



FORMULA

Each 100 ml contains:

Cypermethrin 20 g

Excipients q.s. 100 ml

DESCRIPTION

Concentrated rural insecticide. Provides immediate control and residual effect against insects. Bicherón 50NF is formulated with cypermethrin for insect control in general rural facilities and gardens.

SPECIES

Cattle, sheep, pigs, goats, horses, and dogs.

INDICATIONS

For the control of insects that affect livestock and domestic animals: ticks, house flies, stable flies, mites, horseflies, mosquitoes, lice, fleas, and other insects commonly found in parks, poultry houses, stables, stalls, dairies, livestock sheds, pigpens, and general rural facilities.

ADMINISTRATION ROUTE

Topical, applied as a fine mist to cover surfaces or animals without soaking them, using spraying equipment, mist sprayers, foggers, or fumigators.

DOSAGE

It is recommended to apply a fine mist of the emulsion covering the area to be treated.

In environmental use: approximately 1 liter of emulsion per 5 m².

For animals, the recommended dosage is:

Cattle and horses: 2 to 5 L

Sheep and goats: 2 L (depending on hair or wool length)

Pigs: 1 to 2 L

Dogs: 100 to 200 ml (depending on size)

PREPARATION: Dilute at a ratio of 1:1000 (1 ml per liter of water), stirring gently. Once added, it forms a quick and stable emulsion. It is recommended to use the emulsion within 24 hours of preparation.

PRESENTATIONS

Dropper bottles of 50 and 100 ml.

PRECAUTIONS

Although the product has low toxicity, it is recommended to wear gloves and a mask during handling.

For outdoor use, apply downwind.

For indoor use, begin application from the farthest point from the exit.

Do not eat, drink, or smoke during handling.

Do not store or transport with food.

Do not dispose of in bodies of water, lakes, or ponds, due to its potential toxicity for poikilothermic species (cold-blooded animals).

Do not spray in areas inhabited by cold-blooded animals.

Not recommended for use on puppies.

In case of accidental ingestion, do not induce vomiting or give fatty foods (like milk). Treat with activated charcoal or saline purgatives.

In case of poisoning, seek medical attention.

CONTRAINDICATIONS

Not recommended for use on Oriental dog breeds, as it may cause respiratory and neurological syndromes that are difficult to treat.

WITHDRAWAL PERIOD

A withdrawal period of 72 hours is recommended for meat, eggs, and milk.

STORAGE

Store between 5°C and 25°C.