

Bicherón Curabicheras Plata Forte



FORMULA

Content every 100 ml:

DDVP (2,2-dichlorovinyl-dimethyl phosphate).... 2,00 g

Cypermethrin ((2,2-dichlorovinyl) - 2,2-dimethylcyclopropane

Carboxylate of (RS) - cyano-3- Phenoxybenzyl)... 0,40 g

Silver Sulfadiazine 0,10 g

Diluent and excipients q.s.f.

DESCRIPTION

Bicherón Curabicheras Plata Forte is a product to control myiasis. It has larvicidal, antimicrobial and repellent effect, promoting wound healing.

TARGET SPECIES

Bovine cattle, equine, ovine, goats, swine and canine.

APPLICATION

Shake before using.

Spray 10 cm away on the wound and its edges. It must be applied directly to the wound or the affected area trying to cover it completely.

INDICATIONS

Larvae of *Cochliomyia hominivorax*; *Lucila* sp.; *Calliphora* sp.; *Callitroga* sp. and *Sarcophaga* sp. Gram-positive and Gram-negative bacteria sensitive to sulfadiazine.

DOSAGE

Treatment and prevention of myiasis: a single application is usually effective depending on the wound severity and extent and the environmental conditions.

The interval between applications and the treatment duration can be modified, based on the prescribing veterinarian judgment.

CONTRAINDICATIONS

The product has no contraindications if the usage recommendations are respected.

For external local use only.

WARNINGS AND PRECAUTIONS

Toxic Product Class II (Moderately Toxic). Do not smoke, eat or drink, during use. Keep this medicine out of reach of children and pets. As it is toxic to fish, do not contaminate watercourses. Avoid skin and mucous membrane contact. Wash your hands after each use.

Flammable substance do not burn tin or expose it to temperatures over 50 °C

Poisoning and animal overdose:

In case of intoxication, the symptoms are anxiety, nausea, vomiting, diarrhea, sweating, myosis. The antidote to control the symptoms caused by organophosphate (Dichlorvos) is atropine sulfate at 1 ‰.

There is not specific antidote for cypermethrin.

Poisoning in humans:

The product is toxic to men. In case of accidental intoxication, symptoms are principally characterized by changes in consciousness, muscle weakness, and excessive secretory activity.

Initial treatment should focus on ensuring the maintenance of patent airway and the adequate cardiovascular function.

Adequate decontamination should be carried out according to the route of entry of the toxic, either with a body bath, hair washing, clothing removal, use of activated charcoal at conventional and cathartic doses.

Atropine sulfate is used as the specific antidote for Dichlorvos.

Call the doctor immediately.

Keep out of reach of children.

National Poison Centre (Argentina): 0800-333-0160.

RESTRICTIONS ON USE

Three days must pass between the last treatment and the destination of the animals and milk intended for human consumption.

In case of canine use, avoid the licking of the product with the Elizabethan collar.

PRESENTATION

Net Content: 260g/440 ml

STORAGE

The storage temperature should be between 15 - 30 °C.

The ozone layer is not affected by the aerosol gas.