

Dexametasona Proagro

Dexamethasone 0,2%



FORMULA

Every 100 ml contains:

Dexamethasone 21 phosphate 0,2 g

Vehicle q.s.

DESCRIPTION

Fast-acting corticosteroid anti-inflammatory in ready-to-use injectable solution.

TARGET SPECIES

Bovine cattle, ovine, swine, equine, canine and feline.

INDICATIONS

Inflammatory processes of diverse etiology, ketosis, supportive therapy in fallen cows, inducer of labour in bovines, hypersensitivity and allergies.

ROUTES OF ADMINISTRATION

Intramuscular (IM), intravenous (IV) and intra-synovial routes.

DOSAGE

Bovine cattle: 2,5 to 10 ml

Equine: 2 to 2,5 ml

Ovine: 1,25 to 10 ml

Swine: 0,5 to 1 ml

Dogs and Cats: 0,5 to 1 ml every 10 kg. b.w

PRESENTATIONS

Vial containing 10 and 20 ml.

WARNINGS

It is recommended not to apply to pregnant females, except to professional indication for induction of labour in cattle and sheep.

In extended treatments, the recovery time in the inhibition of hypothalamic–pituitary–adrenal axis should be taken into account, because its function will slowly normalize when the treatment will be interrupted.

CONTRAINDICATIONS

The use of systemic corticosteroids is contraindicated in fungal, viral and bacterial infections in which the corresponding antimicrobial therapy has not been established.

Do not administer in animals suffering from active tuberculosis, severe congestive heart failure, severe arterial hypertension, peptic or duodenal ulcer, epilepsy, osteoporosis, uremic syndrome, diabetes, glaucoma, cataracts, herpetic keratitis, pregnancy and in immunosuppressed animals.

It is suggested not to vaccinate animals that are under prolonged treatment with corticosteroids, or if necessary, they should be revaccinated once it is finished.

Do not perform intra-synovial administration if sepsis is observed in or around the joint, if there is the possibility of bone fracture or damaged cartilage; if there are degenerative bone lesions or osteochondrosis.

WITHDRAWAL PERIOD

Do not slaughter animals for human consumption until 5 days after the last treatment, nor use the milk intended for human consumption or industrialization until 72 hours after the treatment.

STORAGE

The storage temperature should be between 5 and 25 °C.