


Cefalexin Plus 205

Cefalexin 18%



ACTIVE SUBSTANCE

Cefalexin.....180 mg

Excipients q.s.1 ml

SPECIES

Canine and feline

USE

Treatment of infections of the respiratory system, urinary tract, skin, soft tissues and bone and joints diseases caused by cefalexin sensitive organisms.

ADMINISTRATION ROUTE

Parenteral administration: intramuscular or subcutaneous injection.

DOSAGE

Canines and felines: 10 mg/Kg of weight. In both cases, once a day during a period of no less than 5 days. This product can be used in pregnant dogs after 5 days.

DOSAGE TABLE

Animal	Weight	Dose	Suspension Ml
Feline	4.5 kg.	10 mg./kg.	0.25 ml.
Canine	4.5 - 9 kg.	10 mg./kg.	0.25 - 0.5 ml.
	9 - 27 kg.		0.5 - 1.5 ml.
	27 - 54 kg.		1.5 - 3 ml.

RESTRICTION PERIOD

Animals for slaughter: 4 days since last treatment.
Without restriction period in case of bovine milk.

CONTRAINDICATIONS

Do not use this product on animal with kidney failure and sensitivity to the antibiotic. Use dry syringes to avoid inactivation of the remaining cefalexin in the vial.

HOW SUPPLIED

50 ml vials.