

Cefatidina

Cephalexine 500 mg

Ranitidine 40 mg



FORMULA

Cephalexine..... 500 mg

Ranitidine..... 40 mg

Excipients q.s.ad..... 750 mg

PRESENTATION

1 blister with 10 tablets of 750 mg each.

SPECIES

For canine and feline.

INDICATIONS

Wide spectrum antimicrobial associated with an antiulcer a and an acid stomach inhibitor.

For the treatment of the respiratory system, urinary track, skin, gastrointestinal system, soft tissues infections as well as bones and articulations diseases.

DOSAGE

Canines: 1 tablet every 20 kg of weight

Felines: ¼ tablet every 5 kg of weight

Every 8 or 12 hs or at the discretion of the attending professional

CONTRAINDICATIONS AND WARNING

Do not administrate to animals suffering from a renal or hepatic insufficiency.

Do not administrate to animals with sensitivity to the antibiotic or to beta lactamics

The use of this product may produce agranulocytosis sporadically, heart arrythmias, excitability and emesis.

Do not take concomitantly with other antiacids.

Keep out of the reach of the children and store at a temperature between 5 and 25º C.

TREATMEN DURATION

The treatment can be repeates 2 or 3 times a day for 5 days according to the patients evolution and according to the proffesionals judgment.

In case of superficial pyoderma gangrenosum the treatment must continue for 7 to 10 days after the clinical symptoms disappear.

In serious cases of pyoderma the treatment must continue for 14 to 21 days after the clinical symptoms desappear.

In case of acute infections the dose can be administered more frequently.